

Saturating the Bound on the Scale of Fermion Mass Generation

R. Sekhar Chivukula*

Department of Physics, Boston University,
590 Commonwealth Ave., Boston MA 02215

July 16, 1998

Abstract

Recently, Jäger and Willenbrock have shown that the Appelquist and Chanowitz bound on the scale of top-quark mass generation can formally be saturated at tree-level in a particular limit of a two-Higgs doublet model. In this note I present an alternate derivation of their result. I perform a coupled channel analysis for $f\bar{f} \rightarrow V_L V_L$ and $V_L V_L \rightarrow V_L V_L$ scattering and derive the conditions on the parameters required for $f\bar{f} \rightarrow V_L V_L$ scattering to be relevant to unitarity. I also show that it is not possible to saturate the bound on fermion mass generation in a two-Higgs model while maintaining tree-level unitarity in Higgs scattering at high energies.

*e-mail addresses: sekhar@bu.edu

Appelquist and Chanowitz [1] derived an upper bound on the scale of fermion mass generation by examining the inelastic scattering amplitude $f\bar{f} \rightarrow V_L V_L$, where V_L denotes longitudinally polarized W or Z gauge bosons. In the absence of the Higgs boson, or another dynamics responsible for generating fermion mass, this amplitude grows with increasing center-of-mass energy. This tree-level amplitude would ultimately violate unitarity at a sufficiently high energy Λ_f . Therefore, one concludes that the scale associated with fermion mass generation is bounded by Λ_f . The strictest bound for a fermion with mass m_f is obtained from the spin-zero, weak-isosinglet, color-singlet amplitude [2]

$$\Lambda_f < \frac{8\pi v^2}{\sqrt{3N_c}m_f}, \quad (1.1)$$

where $v = 246$ GeV and N_c is the number of colors (3 for quarks and 1 for leptons). The strongest bound occurs for the top quark. With $m_t \approx 175$ GeV, the upper bound (1.1) on the scale of top mass generation is $\Lambda_t \approx 3$ TeV.

On the other hand, in the absence of a Higgs boson or some other dynamics responsible for electroweak gauge-boson mass generation, the elastic scattering amplitude $V_L V_L \rightarrow V_L V_L$ grows quadratically with center-of-mass energy. Therefore, one concludes that the scale associated with gauge boson mass generation is bounded by a scale Λ_{EW} where the elastic scattering amplitude would violate unitarity. The strictest bound is obtained from the spin-zero, weak-isosinglet scattering amplitude [3, 4, 2]

$$\Lambda_{EW} < \sqrt{8\pi}v \approx 1.2 \text{ TeV}. \quad (1.2)$$

Since Λ_{EW} is less than Λ_t , it is not clear that the bound (1.1) is relevant [1, 5]. The physics responsible for unitarizing the elastic gauge boson scattering amplitude may unitarize the inelastic $t\bar{t} \rightarrow V_L V_L$ amplitude, as happens in the standard one-doublet Higgs model.

Recently, Jäger and Willenbrock [6] have shown that the Appelquist and Chanowitz [1] bound (1.1) on the scale of top quark mass generation can formally be saturated at tree-level in a particular limit of a two-Higgs doublet model. In this note I present an alternate derivation of their result. I perform a coupled channel analysis for $f\bar{f} \rightarrow V_L V_L$ and $V_L V_L \rightarrow V_L V_L$ scattering and derive the conditions on the parameters required for $f\bar{f} \rightarrow V_L V_L$ scattering to be relevant to unitarity. I also show that it is not possible to saturate the bound on fermion mass generation in a two-Higgs model while maintaining tree-level unitarity in Higgs scattering at high energies.

Consider the general potential in a two-Higgs model written in the form [7]

$$\begin{aligned} V(\phi_1, \phi_2) &= \lambda_1(\phi_1^\dagger \phi_1 - v_1^2/2)^2 + \lambda_2(\phi_2^\dagger \phi_2 - v_2^2/2)^2 \\ &+ \lambda_3 \left[(\phi_1^\dagger \phi_1 - v_1^2/2) + (\phi_2^\dagger \phi_2 - v_2^2/2) \right]^2 \\ &+ \lambda_4 \left[(\phi_1^\dagger \phi_1)(\phi_2^\dagger \phi_2) - (\phi_1^\dagger \phi_2)(\phi_2^\dagger \phi_1) \right] \end{aligned}$$

$$\begin{aligned}
& + \lambda_5 \left[\text{Re}(\phi_1^\dagger \phi_2) - v_1 v_2 \cos \xi/2 \right]^2 \\
& + \lambda_6 \left[\text{Im}(\phi_1^\dagger \phi_2) - v_1 v_2 \sin \xi/2 \right]^2 ,
\end{aligned} \tag{1.3}$$

where ϕ_1 and ϕ_2 are weak doublet scalar fields with hypercharge $+1/2$. For simplicity, in the following we will set $\xi = 0$. In order to obtain the correct gauge boson masses, we must require that

$$v_1^2 + v_2^2 = v^2 \approx (246 \text{ GeV})^2 . \tag{1.4}$$

We also impose a softly broken discrete symmetry under which ϕ_1 and the right-handed down-quark and charged-lepton fields change sign. This symmetry eliminates an extra terms which would otherwise have been present in (1.3) and it insures that only ϕ_2 contributes to up-quark masses in general and the top-quark mass in particular.

In this language, the limit considered by Jäger and Willenbrock [6] corresponds to $\lambda_5 \rightarrow \infty$ [8], λ_i (where $i = 1 - 4$ or 6) small, and $v_2 \ll v_1 \approx v$. Note that this is a non-decoupling limit [5, 6] in that a *dimensionless* coupling, λ_5 , is taken large instead of a dimensionful one [9].

The vacuum expectation values (vevs) and neutral scalar fields can be written

$$\phi_1 \rightarrow \begin{pmatrix} 0 \\ (h_1 + v_1)/\sqrt{2} \end{pmatrix} \quad \& \quad \phi_2 \rightarrow \begin{pmatrix} 0 \\ (h_2 + v_2)/\sqrt{2} \end{pmatrix} . \tag{1.5}$$

In the limit considered, the dominant contributions to the neutral scalar masses come from the λ_5 term above, which gives:

$$\frac{\lambda_5}{4} [v_2 h_1 + v_1 h_2 + h_1 h_2]^2 . \tag{1.6}$$

From this we immediately see that the combination

$$H = \frac{v_2 h_1 + v_1 h_2}{\sqrt{v_1^2 + v_2^2}} \tag{1.7}$$

has approximately the mass

$$m_H^2 = \frac{\lambda_5 v^2}{2} . \tag{1.8}$$

In contrast, the orthogonal combination

$$h = \frac{-v_1 h_1 + v_2 h_2}{\sqrt{v_1^2 + v_2^2}} \tag{1.9}$$

has mass

$$m_h^2 = \mathcal{O}(\lambda_i v^2) \tag{1.10}$$

with $i = 1-4$ or 6 , and remains light. In the two-Higgs notation employed in [6, 10], these relations may be written

$$\cos \alpha \approx \sin \beta \approx \frac{v_2}{v} , \quad (1.11)$$

and

$$\sin \alpha \approx \cos \beta \approx \frac{v_1}{v} . \quad (1.12)$$

In the limit $\lambda_5 \rightarrow \infty$ and $v_2 \ll v_1$, one can easily verify that the corrections to the expressions given above for the neutral scalar masses and mixings are suppressed by λ_i/λ_5 , v_2/v , or both.

Recalling that only ϕ_2 couples to the top quark, we find the couplings of the neutral scalar fields to the top quark are

$$\frac{\sqrt{2}m_t}{v} \bar{t}t \left(\frac{\cos \alpha}{\sin \beta} h + \frac{\sin \alpha}{\sin \beta} H \right) . \quad (1.13)$$

Note that the coupling of h to the top-quark is approximately equal to that of the standard model Higgs, while the coupling of the H is enhanced by $v_1/v_2 \gg 1$. The couplings of the neutral scalars to W and Z gauge boson pairs is

$$\frac{2}{v} (\sin(\beta - \alpha) h + \cos(\beta - \alpha) H) \left(M_W^2 W^{\mu+} W_\mu^- + \frac{1}{2} M_Z^2 Z^\mu Z_\mu \right) . \quad (1.14)$$

From eqns. (1.11) and (1.12) above, we calculate

$$\sin(\beta - \alpha) \approx \frac{v_2^2 - v_1^2}{v^2} = -1 + \mathcal{O} \left(\frac{v_2^2}{v^2} \right) , \quad (1.15)$$

and

$$\cos(\beta - \alpha) \approx \frac{2v_1 v_2}{v^2} = \frac{2v_2}{v} + \mathcal{O} \left(\frac{v_2^3}{v^3} \right) . \quad (1.16)$$

Next consider scattering amplitudes in the range of energies $m_{W,Z}, m_h, m_t \ll \sqrt{s} \ll m_H$. $V_L V_L$ elastic scattering is largely unitarized by h exchange. The properly normalized tree-level, spin-zero, weak-isosinglet amplitude is

$$\mathcal{A}_{el}(V_L V_L \rightarrow V_L V_L)|_{\sqrt{s} \ll m_H} = \frac{s}{16\pi v^2} (1 - \sin^2(\beta - \alpha)) \approx \frac{s v_2^2}{4\pi v^4} , \quad (1.17)$$

where s is the center-of-mass energy squared.

The scattering amplitude for $t\bar{t} \rightarrow V_L V_L$ grows in this region since $\sqrt{s} < m_H$. The tree-level spin-zero, weak-isosinglet, color singlet amplitude is [2]

$$\mathcal{A}_{inel}(t\bar{t} \rightarrow V_L V_L)|_{\sqrt{s} \ll m_H} = \frac{\sqrt{3N_c} \sqrt{s} m_t}{16\pi v^2} \left(1 - \frac{\cos \alpha}{\sin \beta} \sin(\beta - \alpha) \right) \approx \frac{\sqrt{3N_c} \sqrt{s} m_t}{8\pi v^2} . \quad (1.18)$$

The contribution proportional to $\sin(\beta - \alpha)$ in the amplitude above arises from h -exchange, and while it is equal in magnitude to the contribution from Higgs-exchange in the standard model it has the *opposite* sign!

Finally, the leading contributions to $t\bar{t}$ elastic scattering in this energy regime comes from Z - and h -exchange. As the coupling of the h to $t\bar{t}$ is approximately the same as that of the standard model Higgs, *cf* eqn. (1.13), this amplitude is negligible.

In order to judge the relative importance of $\mathcal{A}_{el}$ and $\mathcal{A}_{inel}$, we perform a coupled-channel analysis of the spin-zero, weak-isosinglet, color-singlet $V_L V_L$ and $t\bar{t}$ states. At tree-level, the amplitudes for $t\bar{t} \rightarrow V_L V_L$ and the reverse process $V_L V_L \rightarrow t\bar{t}$ are real and, therefore, equal. To perform the coupled-channel analysis, we consider the scattering matrix

$$\begin{pmatrix} \mathcal{A}_t \approx 0 & \mathcal{A}_{inel} \\ \mathcal{A}_{inel} & \mathcal{A}_{el} \end{pmatrix}. \quad (1.19)$$

Unitarity requires that the real part of the largest eigenvalue of this matrix be less than one-half [11]. This yields the constraint

$$2\mathcal{A}_{el} + 4\mathcal{A}_{inel}^2 \lesssim 1. \quad (1.20)$$

From this we see that the inelastic amplitude dominates the unitarity constraints in this regime if $2\mathcal{A}_{inel}^2 > \mathcal{A}_{el}$. (Note that, the contributions to overall unitarity from *both* processes scale like s .) From eqns. (1.17) and (1.18) we see this occurs only if

$$v_2 \leq \sqrt{\frac{3N_c}{8\pi}} m_t \approx 105 \text{ GeV}. \quad (1.21)$$

Can v_2 be this small in the two-Higgs doublet model? Consider scattering at high-energies, $\sqrt{s} \gg m_H$. In this region, H -exchange contributes significantly to $t\bar{t}$ elastic scattering. The contribution to the spin-zero color singlet amplitude coming from H exchange is [12]

$$\mathcal{A}_t(t\bar{t} \rightarrow t\bar{t})|_{\sqrt{s} \gg m_H} = -\frac{N_c m_t^2}{16\pi v_2^2}. \quad (1.22)$$

Unitarity implies the absolute value of this amplitude must be less than one-half. This yields the bound

$$v_2 \geq \sqrt{\frac{N_c}{8\pi}} m_t \approx 60 \text{ GeV}. \quad (1.23)$$

Comparing eqns. (1.21) and (1.23), we see that one can consistently arrange for the inelastic amplitude for $t\bar{t} \rightarrow V_L V_L$ to dominate over the elastic amplitude for $V_L V_L \rightarrow V_L V_L$ without violating unitarity in elastic $t\bar{t}$ scattering at high energies.

In order to determine whether one can saturate the bound on top-quark mass generation, however, one must see if the H -boson mass can be made as large as

the bound in eqn. (1.1). That is, from (1.8), we must ask how large λ_5 can be. Consider the tree-level amplitude for spin-zero $hH \rightarrow hH$ scattering. From (1.6), we calculate

$$\mathcal{A}_{hH}(hH \rightarrow hH)|_{\sqrt{s} \gg m_H} = -\frac{\lambda_5}{16\pi} . \quad (1.24)$$

Requiring that the real part of this amplitude not exceed one half (in absolute value), we find $\lambda_5 \leq 8\pi$ and hence, from (1.8),

$$m_H \leq \sqrt{4\pi}v \approx 870 \text{ GeV} . \quad (1.25)$$

Since the bound (1.25) is much less than 3 TeV, we conclude that it is not possible to saturate the bound on top-quark mass generation in the two-Higgs model. This conclusion is consistent with that obtained by considering the triviality of the model [13].

In summary, while it is possible in the two-Higgs model to arrange for the inelastic scattering amplitude $t\bar{t} \rightarrow V_L V_L$ to dominate over the elastic $V_L V_L \rightarrow V_L V_L$ amplitude, one cannot saturate the bound on top-quark mass generation while maintaining unitarity in Higgs scattering at high energies. The situation is analogous to that of trying to saturate the scale Λ_{EW} [3, 4, 2] in the standard one-doublet Higgs model. In that case as well, Higgs scattering [3] and triviality [14] preclude making the Higgs boson as heavy as Λ_{EW} .

Finally, we note that the lower bound on v_2 (1.23) is approximately saturated by the “top-Higgs” in topcolor-assisted technicolor models [15]. However, in the simplest version of this model the top-Higgs mass is proportional to v_2 , unlike the relation found in eqn. (1.8). It is interesting to speculate whether the bound (1.1) could be saturated in a variant topcolor-assisted technicolor model.

Acknowledgments

I thank Tom Appelquist, Bogdan Dobrescu, Howard Georgi, Tao Han, Elizabeth Simmons, and Scott Willenbrock for discussions and comments on the manuscript, and the Aspen Center for Physics for its hospitality while this work was completed. *This work was supported in part by the Department of Energy under grant DE-FG02-91ER40676.*

References

- [1] T. Appelquist and M. S. Chanowitz, Phys. Rev. Lett. **59**, 2405 (1987).
- [2] W. Marciano, G. Valencia, and S. Willenbrock, Phys. Rev. **D40**, 1725 (1989).
- [3] B. W. Lee, C. Quigg, and H. B. Thacker, Phys. Rev. **D16**, 1519 (1977).
- [4] M. S. Chanowitz and M. K. Gaillard, Nucl. Phys. **B261**, 379 (1985).

- [5] M. Golden, Phys. Lett. **B338**, 295 (1994), hep-ph/9408272.
- [6] S. Jaeger and S. Willenbrock, Unitarity and the scale of fermion mass generation, hep-ph/9806286, 1998.
- [7] H. Georgi, Hadronic J. **1**, 155 (1978).
- [8] H. E. Haber, (1995), hep-ph/9505240.
- [9] T. Appelquist and J. Carazzone, Phys. Rev. **D11**, 2856 (1975).
- [10] J. F. Gunion, H. E. Haber, G. L. Kane, and S. Dawson, *The Higgs Hunter's Guide* (Addison-Wesley, 1990).
- [11] M. Luscher and P. Weisz, Phys. Lett. **212B**, 472 (1988).
- [12] M. S. Chanowitz, M. A. Furman, and I. Hinchliffe, Nucl. Phys. **B153**, 402 (1979).
- [13] D. Kominis and R. S. Chivukula, Phys. Lett. **B304**, 152 (1993), hep-ph/9301222.
- [14] R. Dashen and H. Neuberger, Phys. Rev. Lett. **50**, 1897 (1983).
- [15] C. T. Hill, Phys. Lett. **B345**, 483 (1995), hep-ph/9411426.